

M2REG-DEFINT-2425-ASM-SET 6-MATH

建議題解

1. (a) 設 $u = a - x$ 。則 $du = -dx$ 。 1M

$$\begin{aligned}\int_0^a f(a-x) dx &= -\int_a^0 f(u) du \\ &= \int_0^a f(u) du & 1M \\ &= \int_0^a f(x) dx & 1\end{aligned}$$

可得 $\int_0^a f(x) dx = \int_0^a f(a-x) dx$ 。

$$\begin{aligned}\int_0^a f(x) dx &= \frac{1}{2} \left[\int_0^a f(x) dx + \int_0^a f(a-x) dx \right] \\ &= \frac{1}{2} \int_0^a [f(x) + f(a-x)] dx & 1\end{aligned}$$

(b) $\int_0^3 \frac{dx}{e^{3-2x} + 1}$

$$= \frac{1}{2} \int_0^3 \left[\frac{1}{e^{3-2x} + 1} + \frac{1}{e^{3-2(3-x)} + 1} \right] dx \quad 1M$$

$$\begin{aligned}&= \frac{1}{2} \int_0^3 \left[\frac{1}{e^{3-2x} + 1} + \frac{1}{e^{2x-3} + 1} \times \frac{e^{3-2x}}{e^{3-2x}} \right] dx \\ &= \frac{1}{2} \int_0^3 \frac{1 + e^{3-2x}}{e^{3-2x} + 1} dx & 1M\end{aligned}$$

$$\begin{aligned}&= \frac{1}{2} \left[x \right]_0^3 \\ &= \frac{3}{2} & 1A\end{aligned}$$

$$\begin{aligned}
 2. \quad (a) \quad (i) \quad \cos \theta (2 \cos 4\theta - 2 \cos 2\theta + 1) &= 2 \cos \theta \cos 4\theta - 2 \cos \theta \cos 2\theta + \cos \theta \\
 &= (\cos 5\theta + \cos 3\theta) - (\cos 3\theta + \cos \theta) + \cos \theta \\
 &= \cos 5\theta
 \end{aligned}$$

1M

$$\text{因此, } \frac{\cos 5\theta}{\cos \theta} = 2 \cos 4\theta - 2 \cos 2\theta + 1.$$

1

$$(ii) \quad \frac{\cos 5\left(\frac{\pi}{4} - t\right)}{\cos\left(\frac{\pi}{4} - t\right)} = 2 \cos\left(\frac{\pi}{4} - t\right) - 2 \cos 2\left(\frac{\pi}{4} - t\right) + 1$$

$$\frac{\cos \frac{5\pi}{4} \cos 5t + \sin \frac{5\pi}{4} \sin 5t}{\cos \frac{\pi}{4} \cos t + \sin \frac{\pi}{4} \sin t} = -2 \cos 4t - 2 \sin 2t + 1$$

1M

$$\frac{-\frac{\sqrt{2}}{2} \cos 5t - \frac{\sqrt{2}}{2} \sin 5t}{\frac{\sqrt{2}}{2} \cos t + \frac{\sqrt{2}}{2} \sin t} = -2 \cos 4t - 2 \sin 2t + 1$$

$$\frac{\cos 5t + \sin 5t}{\cos t + \sin t} = 2 \cos 4t + 2 \sin 2t - 1$$

1

$$(b) \quad (i) \quad \text{設 } u = a - x. \text{ 則 } du = -dx.$$

1M

$$\begin{aligned}
 \int_0^a f(a-x) dx &= -\int_a^0 f(u) du \\
 &= \int_0^a f(u) du
 \end{aligned}$$

1M

$$= \int_0^a f(x) dx$$

1

$$(ii) \quad \int_0^{\frac{\pi}{2}} \frac{\cos 5x}{\cos x + \sin x} dx = \int_0^{\frac{\pi}{2}} \frac{\cos 5\left(\frac{\pi}{2} - x\right)}{\cos\left(\frac{\pi}{2} - x\right) + \sin\left(\frac{\pi}{2} - x\right)} dx$$

1M

$$= \int_0^{\frac{\pi}{2}} \frac{\sin 5x}{\cos x + \sin x} dx$$

1

$$\begin{aligned}
 (c) \quad &\int_0^{\frac{\pi}{2}} \frac{\cos 5x}{\cos x + \sin x} dx \\
 &= \frac{1}{2} \left[\int_0^{\frac{\pi}{2}} \frac{\cos 5x}{\cos x + \sin x} dx + \int_0^{\frac{\pi}{2}} \frac{\sin 5x}{\cos x + \sin x} dx \right]
 \end{aligned}$$

1M

$$= \frac{1}{2} \int_0^{\frac{\pi}{2}} (2 \cos 4x + 2 \sin 2x - 1) dx$$

1M

$$= \frac{1}{2} \left[\frac{\sin 4x}{2} - \cos 2x - x \right]_0^{\frac{\pi}{2}}$$

$$= 1 - \frac{\pi}{4}$$

1A

3. (a) 設 $u = a - x$ 。則 $du = -dx$ 。 1M

$$\begin{aligned}\int_0^a f(a-x) dx &= -\int_a^0 f(u) du \\ &= \int_0^a f(u) du & 1M \\ &= \int_0^a f(x) dx & 1\end{aligned}$$

$$\begin{aligned}\text{(b)} \quad \int_0^{\frac{\pi}{4}} \ln(\cos x) dx &= \int_0^{\frac{\pi}{4}} \ln\left[\cos\left(\frac{\pi}{4} - x\right)\right] dx & 1M \\ &= \int_0^{\frac{\pi}{4}} \ln\left(\cos\frac{\pi}{4}\cos x + \sin\frac{\pi}{4}\sin x\right) dx & 1M \\ &= \int_0^{\frac{\pi}{4}} \ln\frac{\cos x + \sin x}{\sqrt{2}} dx & 1\end{aligned}$$

$$\begin{aligned}\text{(c)} \quad \int_0^{\frac{\pi}{4}} \ln(\cos x) dx &= \int_0^{\frac{\pi}{4}} \ln\frac{\cos x + \sin x}{\sqrt{2}} dx \\ &= \int_0^{\frac{\pi}{4}} \ln(\cos x + \sin x) dx - \int_0^{\frac{\pi}{4}} \sqrt{2} dx & 1M\end{aligned}$$

$$\int_0^{\frac{\pi}{4}} \ln\frac{\cos x}{\cos x + \sin x} dx = -\ln\sqrt{2}\left[x\right]_0^{\frac{\pi}{4}} \quad 1M$$

$$\int_0^{\frac{\pi}{4}} \frac{1}{1 + \tan x} dx = -\frac{\pi \ln 2}{8} \quad 1$$

$$\text{(d)} \quad \text{留意 } \frac{d}{dx} [\ln(1 + \tan x)] = \frac{\sec^2 x}{1 + \tan x}。$$

$$\int_0^{\frac{\pi}{4}} \frac{x \sec^2 x}{1 + \tan x} dx = \left[x \ln(1 + \tan x)\right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx \quad 1M$$

$$\begin{aligned}&= \frac{\pi \ln 2}{4} + \int_0^{\frac{\pi}{4}} \ln\frac{1}{1 + \tan x} dx \\ &= \frac{\pi \ln 2}{4} + \left(-\frac{\pi \ln 2}{8}\right) & 1M\end{aligned}$$

$$= \frac{\pi \ln 2}{8} \quad 1A$$

4. (a) 設 $u = \pi - x$ 。則 $du = -dx$ 。 1M

$$\begin{aligned} \int_{\frac{\pi}{2}}^{\pi} e^{\frac{1}{1+\sin x}} f(x) dx &= - \int_{\frac{\pi}{2}}^0 e^{\frac{1}{1+\sin(\pi-u)}} f(\pi-u) du \\ &= \int_0^{\frac{\pi}{2}} e^{\frac{1}{1+\sin u}} [-f(u)] du \end{aligned} \quad 1M$$

$$= - \int_0^{\frac{\pi}{2}} e^{\frac{1}{1+\sin x}} f(x) dx \quad 1$$

$$\begin{aligned} \int_0^{\pi} e^{\frac{1}{1+\sin x}} f(x) dx &= \int_0^{\frac{\pi}{2}} e^{\frac{1}{1+\sin x}} f(x) dx + \int_{\frac{\pi}{2}}^{\pi} e^{\frac{1}{1+\sin x}} f(x) dx \\ &= \int_0^{\frac{\pi}{2}} e^{\frac{1}{1+\sin x}} f(x) dx - \int_0^{\frac{\pi}{2}} e^{\frac{1}{1+\sin x}} f(x) dx \\ &= 0 \end{aligned} \quad 1$$

(b) 設 $v = \pi - x$ 。則 $dv = -dx$ 。 1M

$$\begin{aligned} \int_0^{\pi} \sin x g(x) dx &= \int_0^{\frac{\pi}{2}} \sin x g(x) dx + \int_{\frac{\pi}{2}}^{\pi} \sin x g(x) dx \\ &= \int_0^{\frac{\pi}{2}} \sin x g(x) dx - \int_{\frac{\pi}{2}}^0 \sin(\pi-u) g(\pi-u) du \\ &= \int_0^{\frac{\pi}{2}} \sin x g(x) dx + \int_0^{\frac{\pi}{2}} \sin u g(u) du \end{aligned} \quad 1M$$

$$\begin{aligned} &= \int_0^{\frac{\pi}{2}} \sin x g(x) dx + \int_0^{\frac{\pi}{2}} \sin x g(x) dx \\ &= 2 \int_0^{\frac{\pi}{2}} \sin x g(x) dx \end{aligned} \quad 1$$

(c) (i) 代 $f(x) = \cos x \sin x$ 。

$$f(\pi - x) = \cos(\pi - x) \sin(\pi - x) = -\cos x \sin x = -f(x) \quad 1M$$

$$\int_0^{\pi} e^{\frac{1}{1+\sin x}} \cos x \sin x dx = 0 \quad 1A$$

設 $w = \cos x$ 。則 $dw = -\sin x dx$ 。 1M

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \sin x \cos^2 x dx &= - \int_1^0 u^2 du \\ &= -\frac{1}{3} \left[u^3 \right]_1^0 \\ &= \frac{1}{3} \end{aligned} \quad \begin{array}{l} 1A \\ 1A \end{array}$$

$$\begin{aligned} \text{(ii)} \quad \int_0^{\pi} \sin 2x (\cos x + e^{\frac{1}{1+\sin x}}) dx &= \int_0^{\pi} (\sin 2x \cos x + e^{\frac{1}{1+\sin x}} \sin 2x) dx \\ &= 2 \int_0^{\pi} \sin x \cos^2 x dx + 2 \int_0^{\pi} e^{\frac{1}{1+\sin x}} \sin x \cos x dx \\ &= 2 \int_0^{\pi} \sin x \cos^2 x dx \end{aligned} \quad 1M$$

$$\text{代 } g(x) = \cos^2 x \circ g(\pi - x) = \cos^2(\pi - x) = \cos^2 x = g(x) \circ \quad 1\text{M}$$

$$\begin{aligned} \int_0^{\pi} \sin 2x (\cos x + e^{\frac{1}{1+\sin x}}) dx &= 2 \int_0^{\pi} \sin x \cos^2 x dx \\ &= 4 \int_0^{\frac{\pi}{2}} \sin x \cos^2 x dx \\ &= 4 \left(\frac{1}{3} \right) \\ &= \frac{4}{3} \end{aligned} \quad 1\text{A}$$

$$5. \quad (\text{a}) \quad \text{設 } u = x^3 \circ \text{則 } du = 3x^2 dx \circ \quad 1\text{M}$$

$$\begin{aligned} \int x^2 \sin x^3 dx &= \frac{1}{3} \int \sin u du \\ &= -\frac{\cos u}{3} + \text{常數} \\ &= -\frac{\cos x^3}{3} + \text{常數} \end{aligned} \quad 1\text{A}$$

$$(i) \quad \text{設 } u = a + b - x \circ \text{則 } du = -dx \circ \quad 1\text{M}$$

$$\begin{aligned} &\int_a^b \frac{f(x)}{f(a+b-x) + f(x)} dx \\ &= - \int_b^a \frac{f(a+b-u)}{f(u) + f(a+b-u)} du \\ &= \int_a^b \frac{f(a+b-u)}{f(u) + f(a+b-u)} du \end{aligned} \quad 1\text{M}$$

$$= \int_a^b \frac{f(a+b-x)}{f(x) + f(a+b-x)} dx \quad 1\text{A}$$

$$\begin{aligned} &\int_a^b \frac{f(x)}{f(a+b-x) + f(x)} dx \\ &= \frac{1}{2} \left[\int_a^b \frac{f(x)}{f(a+b-x) + f(x)} dx + \int_a^b \frac{f(a+b-x)}{f(a+b-x) + f(x)} dx \right] \end{aligned} \quad 1\text{M}$$

$$= \frac{1}{2} \int_a^b dx \quad 1\text{A}$$

$$\begin{aligned} &= \frac{1}{2} \left[x \right]_a^b \\ &= \frac{b-a}{2} \end{aligned} \quad 1$$

$$(ii) \quad \text{設 } u = x^3 \circ \text{則 } du = 3x^2 dx \circ \quad 1\text{M}$$

$$\begin{aligned} &\int_{\sqrt[3]{\log 5}}^{\sqrt[3]{\log 6}} \frac{x^2 \sin x^3}{\sin(\log 30 - x^3) + \sin x^3} dx \\ &= \frac{1}{3} \int_{\log 5}^{\log 6} \frac{\sin u}{\sin(\log 5 + \log 6 - u) + \sin u} du \end{aligned} \quad 1\text{M}$$

$$= \frac{1}{3} \cdot \frac{\log 6 - \log 5}{2} \quad 1\text{M}$$

$$= \frac{1}{6} \log \frac{6}{5} \quad 1\text{A}$$

6. (a) 設 $u = \pi - x$ 。則 $du = -dx$ 。 1M

$$\begin{aligned}
 \int_0^\pi x f(\sin x) dx &= - \int_\pi^0 (\pi - u) f(\sin(\pi - u)) du \\
 &= \int_0^\pi (\pi - u) f(\sin u) du \\
 &= \pi \int_0^\pi f(\sin u) du - \int_0^\pi u f(\sin u) du \\
 &= \pi \int_0^\pi f(\sin x) dx - \int_0^\pi x f(\sin x) dx \\
 2 \int_0^\pi x f(\sin x) dx &= \pi \int_0^\pi f(\sin x) dx \\
 \int_0^\pi x f(\sin x) dx &= \frac{\pi}{2} \int_0^\pi f(\sin x) dx
 \end{aligned}$$

1M

1

(b) (i) 設 $v = -x$ 。則 $dv = -dx$ 。 1M

$$\begin{aligned}
 \int_{-\pi}^0 \frac{g(x)}{1+e^x} dx &= - \int_\pi^0 \frac{g(-v)}{1+e^{-v}} dv \\
 &= \int_0^\pi \frac{e^v g(v)}{e^v(1+e^{-v})} dv \\
 &= \int_0^\pi \frac{e^v g(v)}{1+e^v} dv \\
 &= \int_0^\pi \frac{e^x g(x)}{1+e^x} dx
 \end{aligned}$$

1

$$\begin{aligned}
 \text{(ii)} \quad \int_{-\pi}^\pi \frac{g(x)}{1+e^x} dx &= \int_{-\pi}^0 \frac{g(x)}{1+e^x} dx + \int_0^\pi \frac{g(x)}{1+e^x} dx \\
 &= \int_0^\pi \frac{e^x g(x)}{1+e^x} dx + \int_0^\pi \frac{g(x)}{1+e^x} dx \\
 &= \int_0^\pi \frac{(1+e^x)g(x)}{1+e^x} dx \\
 &= \int_0^\pi g(x) dx
 \end{aligned}$$

1

(c) 代 $g(x) = \frac{\sqrt{2+\cos x}}{\sqrt{2+\cos x} + \sqrt{2-\cos x}}$ 。 1M

$$g(-x) = \frac{\sqrt{2+\cos(-x)}}{\sqrt{2+\cos(-x)} + \sqrt{2-\cos(-x)}} = \frac{\sqrt{2+\cos x}}{\sqrt{2+\cos x} + \sqrt{2-\cos x}} = g(x)$$

$g(x)$ 為在 $[-\pi, \pi]$ 上的連續偶函數。

$$\begin{aligned}
 &\int_{-\pi}^\pi \frac{\sqrt{2+\cos x}}{(1+e^x)(\sqrt{2+\cos x} + \sqrt{2-\cos x})} dx \\
 &= \int_0^\pi g(x) dx \\
 &= \left[xg(x) \right]_0^\pi - \int_0^\pi xg'(x) dx
 \end{aligned}$$

1M

$$g'(x) = \frac{\frac{-\sin x}{2\sqrt{2+\cos x}}(\sqrt{2+\cos x} + \sqrt{2-\cos x}) - (\sqrt{2+\cos x})\left(\frac{-\sin x}{2\sqrt{2+\cos x}} + \frac{\sin x}{2\sqrt{2-\cos x}}\right)}{(\sqrt{2+\cos x} + \sqrt{2-\cos x})^2} \quad 1M$$

$$= \frac{-\sin x}{2\sqrt{3+\sin^2 x} + (3+\sin^2 x)} \quad 1A$$

取 $f(x) = \frac{-x}{2\sqrt{3+x^2} + 3+x^2}$ 使得它在 $[0, 1]$ 連續及 $g'(x) = f(\sin x)$ 。

$$\begin{aligned} & \int_{-\pi}^{\pi} \frac{\sqrt{2+\cos x}}{(1+e^x)(\sqrt{2+\cos x} + \sqrt{2-\cos x})} dx \\ &= \left[xg(x) \right]_0^{\pi} - \int_0^{\pi} xf(\sin x) dx \\ &= \pi g(\pi) - \frac{\pi}{2} \int_0^{\pi} f(\sin x) dx \quad 1M \end{aligned}$$

$$\begin{aligned} &= \pi g(\pi) - \frac{\pi}{2} \int_0^{\pi} g'(x) dx \\ &= \pi g(\pi) - \frac{\pi}{2} \left[g(x) \right]_0^{\pi} \\ &= \frac{\pi}{2} \left(\frac{\sqrt{2-1}}{\sqrt{2-1} + \sqrt{2-1}} + \frac{\sqrt{2+1}}{\sqrt{2+1} + \sqrt{2-1}} \right) \\ &= \frac{\pi}{2} \quad 1A \end{aligned}$$